

MONIKA BUDZYŃSKA, ALEKSANDRA GRZESIK
and MARIOLA KOT

The generalized Day norm Part II. Applications

ABSTRACT. In this paper we prove that for each $1 < p, \tilde{p} < \infty$, the Banach space $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ can be equivalently renormed in such a way that the Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ is LUR and has a diametrically complete set with empty interior. This result extends the Maluta theorem about existence of such a set in l^2 with the Day norm. We also show that the Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ has the weak fixed point property for nonexpansive mappings.

1. Introduction. Recently E. Maluta presented a reflexive LUR Banach space which contains a diametrically complete set with empty interior [11]. Namely she proved that the Banach space l^2 furnished with the Day norm $\|\cdot\|_L$ is such a Banach space. Next in [3] the authors introduced the generalized Day norm $\|\cdot\|_{\beta,p}$ in c_0 and showed its properties. In this paper applying the generalized Day norm, we prove that for each $1 < p, \tilde{p} < \infty$ in the Banach space $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ there exists an equivalent norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ such that $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ contains diametrically complete set with empty interior. We also show that the Banach spaces $(c_0, \|\cdot\|_{\beta,p})$ and $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ have the weak fixed point property for nonexpansive mappings.

2010 *Mathematics Subject Classification.* 46G20, 47H10, 52A05.

Key words and phrases. Diametrically complete set, Day norm, fixed point, Kadec–Klee property, LUR space, nonexpansive mapping, non-strict Opial property, 1-unconditional Schauder bases.

2. Basic notions and facts. Throughout this paper all Banach spaces are infinite dimensional and real. We will also use the notations, assumptions and facts from [3]. Additionally, let us recall the following definition.

Definition 2.1 ([6] and [8], see also [1], [4] and [5]). Let $(X, \|\cdot\|)$ be a Banach space. We say that $(X, \|\cdot\|)$ has the *Kadec–Klee property* with respect to the weak topology (the Kadec–Klee property, for short) if each sequence $\{x_n\}$ with $\lim_n \|x_n\| = 1$, which converges weakly to a point ξ with $\|\xi\| = 1$, tends strongly to ξ .

It is known that for $1 < p < +\infty$ the Banach space l^p with the standard norm $\|\cdot\|_p$ has the Kadec–Klee property (see for example [4]).

We also need the definition of a diametrically complete set in a Banach space.

Definition 2.2. Let $(X, \|\cdot\|)$ be an infinite dimensional Banach space and let C be a nonempty and non-singleton subset of X . We say that C is a diametrically complete set in X if

$$\begin{aligned} \text{diam}_{\|\cdot\|}(C \cup \{x\}) &= \sup\{\|y - y'\| : y, y' \in C \cup \{x\}\} \\ &> \text{diam}_{\|\cdot\|}(C) = \sup\{\|y - y'\| : y, y' \in C\} \end{aligned}$$

for each $x \in X \setminus C$.

It is obvious that a diametrically complete set has to be bounded, closed and convex.

Next we give two results which establish relations between a diametral property of a set and the interior of a diametrically complete set. First in [14], J. P. Moreno, P. L. Papini and R. R. Phelps proved the following theorem.

Theorem 2.3. *Let $(X, \|\cdot\|)$ be an infinite dimensional Banach space and $C \subset X$ be diametrically complete. If the interior of C is empty, then C is diametral.*

In [12], E. Maluta and P. L. Papini showed the following result.

Theorem 2.4. *Each infinite dimensional and reflexive Banach space $(X, \|\cdot\|)$, which satisfies the non-strict Opial property and lacks normal structure, contains diametrically complete sets whose interior is empty.*

3. The generalized Day norm and renorming of Banach spaces.

In this section we apply the generalized Day norm $\|\cdot\|_{\beta,p}$ in c_0 to renorm a Banach space $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$, where $1 < p, \tilde{p} < +\infty$.

Fix $\alpha \in (0, 1)$ and fix $1 < p, \tilde{p} < \infty$. Next we choose a strictly decreasing positive sequence $\beta = \{\beta_j\}_j$ satisfying the following two conditions

- the series $\sum_{j=1}^{\infty} \beta_j^p$ is convergent,

- there exists a constant $L > 1$ such that for each $n \in \mathbb{N}$

$$\sum_{j=n+1}^{\infty} \beta_j^p \leq L\beta_{n+1}^p.$$

Now we can observe that for each $x = \{x^k\}_k \in l^{\tilde{p}}$, the sequence given by

$$u(x) = \{u^i(x)\}_i = \{\alpha\|x\|_{\tilde{p}}, x^1, x^2, x^2, \dots, x^k, \dots, x^k, \dots\}$$

is an element of c_0 (here the k -th coordinate of x is repeated exactly k times and $\|\cdot\|_{\tilde{p}}$ is the standard norm in $l^{\tilde{p}}$). So we can apply the Day norm $\|\cdot\|_{\beta,p}$ ([3]) to the element $u(x)$ and set

$$\|x\|_{L,\alpha,\beta,p,\tilde{p}} := \|u(x)\|_{\beta,p} = \|D(u(x))\|_p,$$

where $\|\cdot\|_p$ is the standard norm in l^p .

It is easy to note that

$$\|u(x)\|_{c_0} = \max\{\alpha\|x\|_{\tilde{p}}, |x^1|, |x^2|, \dots\},$$

$$\beta_1\alpha\|x\|_{\tilde{p}} \leq \|D(u(x))\|_p = \|x\|_{L,\alpha,\beta,p,\tilde{p}} \leq \left(\sum_{j=1}^{\infty} \beta_j^p\right)^{\frac{1}{p}} \|x\|_{\tilde{p}},$$

and

$$\begin{aligned} \beta_1\|u(x)\|_{c_0} &\leq \|D(u(x))\|_p = \|u(x)\|_{\beta,p} = \|x\|_{L,\alpha,\beta,p,\tilde{p}} \\ &\leq \left(\sum_{j=1}^{\infty} \beta_j^p\right)^{\frac{1}{p}} \|u(x)\|_{c_0} \end{aligned}$$

for each $x \in l^{\tilde{p}}$. Therefore $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ is a norm in $l^{\tilde{p}}$, which is equivalent to the original one.

Remark 3.1. The norm $\|\cdot\|_L$ connected with the Day norm $\|\cdot\|$ was introduced by M. Smith ([16]) and our norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ is a generalization of the norm $\|\cdot\|_L$. In his paper M. Smith proves that $(l^2, \|\cdot\|_L)$ is a reflexive, locally uniformly rotund Banach space that is not uniformly convex in every direction. In [17], M. Smith and B. Turett show that $(l^2, \|\cdot\|_L)$ lacks normal structure.

4. The norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ and the non-strict Opial property. To get the result about the non-strict Opial property of the Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ we modify the Maluta proof of Theorem 3.1 in [11] (see also the proof of Theorem 5.2 in [3]).

Theorem 4.1. *The Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ has the non-strict Opial property.*

Proof. Assume that $\{x_n\}_n = \{\{x_n^i\}_i\}_n \subset l^{\tilde{p}}$ tends weakly to 0 in $l^{\tilde{p}}$ and let $x = \{x^i\}_i \in l^{\tilde{p}} \setminus \{0\}$. Let us take $0 < \epsilon < 1$. By the Opial property of the Banach space $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ there exists $\tilde{n}_0 \in \mathbb{N}$ such that for each $\tilde{n}_0 < n \in \mathbb{N}$ we have

$$\|x_n\|_{\tilde{p}} < \|x_n - x\|_{\tilde{p}} + \epsilon.$$

Next there exists $\tilde{i} \in \mathbb{N}$ such that

$$|x^i| < \epsilon$$

for each $\tilde{i} < i \in \mathbb{N}$. Therefore,

$$|x_n^i| \leq |x_n^i - x^i| + |x^i| < |x_n^i - x^i| + \epsilon$$

for each $\tilde{i} < i \in \mathbb{N}$ and all $n \in \mathbb{N}$.

Now for each $1 \leq i \leq \tilde{i}$ we have either $x^i = 0$ or $x^i \neq 0$. In the second case setting $\eta_i = \min\{\epsilon, \frac{1}{2}|x^i|\}$ and taking into account the weak convergence of $\{x_n\}_n$ to 0, we find $\tilde{n}_i \in \mathbb{N}$ such that

$$|x_n^i| < \eta_i$$

for $\tilde{n}_i < n \in \mathbb{N}$ and hence we obtain

$$\begin{aligned} |x_n^i - x^i| &\geq |x^i| - |x_n^i| \\ &> |x^i| - \eta_i \geq \frac{1}{2}|x^i| \geq \eta_i > |x_n^i|. \end{aligned}$$

In the first case, i.e., $x^i = 0$, we have

$$|x_n^i| = |x_n^i - x^i|$$

for each $n \in \mathbb{N}$.

So we have shown that

$$|x_n^i| \leq |x_n^i - x^i|$$

for each $1 \leq i \leq \tilde{i}$ and all $\max\{\tilde{n}_1, \dots, \tilde{n}_{\tilde{i}}\} < n \in \mathbb{N}$.

Putting together all above inequalities, we get

$$|x_n^i| < |x_n^i - x^i| + \epsilon$$

for each $i \in \mathbb{N}$ and for all $\max\{\tilde{n}_1, \dots, \tilde{n}_{\tilde{i}}\} < n \in \mathbb{N}$.

The above considerations yield the following inequalities

$$(*) \quad |u^i(x_n)| < |u^i(x_n - x)| + \epsilon$$

for each $i \in \mathbb{N}$ and for all $\max\{\tilde{n}_0, \tilde{n}_1, \dots, \tilde{n}_{\tilde{i}}\} < n \in \mathbb{N}$.

Now let us take $\max\{\tilde{n}_0, \tilde{n}_1, \dots, \tilde{n}_{\tilde{i}}\} < n \in \mathbb{N}$. Then using Corollary 2.8 from [3] and applying (*), we obtain

$$\begin{aligned} \|x_n\|_{L,\alpha,\beta,p,\tilde{p}} &= \|u(x_n)\|_{\beta,p} = \|D(u(x_n))\|_{\tilde{p}} \\ &= \left[\sum_{j=1}^{\infty} \left(\beta_j \left| u^{\tau(j,u(x_n))}(x_n) \right| \right)^p \right]^{\frac{1}{p}} \end{aligned}$$

$$\begin{aligned}
&< \left\{ \sum_{j=1}^{\infty} \beta_j \left[\left| u^{\tau(j, u(x_n))}(x_n - x) \right| + \epsilon \right]^p \right\}^{\frac{1}{p}} \\
&\leq \left\{ \sum_{j=1}^{\infty} \beta_j \left[\left| u^{\tau(j, u(x_n - x))}(x_n - x) \right| + \epsilon \right]^p \right\}^{\frac{1}{p}} \\
&\leq \left[\sum_{j=1}^{\infty} \left(\beta_j \left| u^{\tau(j, u(x_n - x))}(x_n - x) \right| \right)^p \right]^{\frac{1}{p}} + \epsilon \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}} \\
&= \|D(u(x_n - x))\|_p + \epsilon \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}} \\
&= \| \|u(x_n - x)\|_{\beta, p} + \epsilon \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}} \\
&= \|x_n - x\|_{L, \alpha, \beta, p, \bar{p}} + \epsilon \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.
\end{aligned}$$

Finally, by passing n to $+\infty$, we get

$$\limsup_n \|x_n\|_{L, \alpha, \beta, p, \bar{p}} \leq \limsup_n \|x_n - x\|_{L, \alpha, \beta, p, \bar{p}} + \epsilon \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}$$

and by arbitrariness of $0 < \epsilon < 1$, we obtain

$$\limsup_n \|x_n\|_{L, \alpha, \beta, p, \bar{p}} \leq \limsup_n \|x_n - x\|_{L, \alpha, \beta, p, \bar{p}}.$$

□

Observe that the Banach space $(l^{\bar{p}}, \| \cdot \|_{L, \alpha, \beta, p, \bar{p}})$ does not have the Opial property as the following example shows.

Example 4.2. Consider $(l^{\bar{p}}, \| \cdot \|_{L, \alpha, \beta, p, \bar{p}})$ with the standard basis $\{e_i\}_i$. Let us take a sequence $\{u_n\}_n = \{e_{n+1}\}_n$. This sequence is weakly convergent to $0 \in c_0$ and for

$$u = \min \left\{ 1, \left(\frac{1}{\alpha^{\bar{p}}} - 1 \right)^{\frac{1}{\bar{p}}} \right\} e_1$$

we have

$$\lim_n \| \|u_n\|_{L, \alpha, \beta, p, \bar{p}} = \lim_n \| \|u_n - u\|_{L, \alpha, \beta, p, \bar{p}} = \left(\sum_{j=1}^{\infty} \beta_j^p \right)^{\frac{1}{p}}.$$

5. The norm $\|\cdot\|_{L,\alpha,\beta,p,\bar{p}}$ and normal structure. The following theorem is valid.

Theorem 5.1. *The Banach space $(l^{\bar{p}}, \|\cdot\|_{L,\alpha,\beta,p,\bar{p}})$ lacks normal structure for $\alpha \leq 2^{-\frac{1}{\bar{p}}}$.*

Proof. As usual in $l^{\bar{p}}$ we have the standard basis $\{e_i\}_i$. The proof is a small modification of the proof due to M. A. Smith and B. Turett ([17]). Observe that for $m_2 > m_1$ we have

$$e_{m_2} - e_{m_1} = \{0, \dots, 0, -1, 0, \dots, 0, 1, 0, \dots\}.$$

Clearly,

$$\alpha \|e_{m_2} - e_{m_1}\|_{\bar{p}} = \alpha 2^{\frac{1}{\bar{p}}} \leq 1$$

and therefore

$$\begin{aligned} \left[\sum_{j=1}^{m_1+m_2} \beta_j^p \right]^{\frac{1}{p}} &\leq \|e_{m_2} - e_{m_1}\|_{L,\alpha,\beta,p,\bar{p}} = \|e_{m_2} - e_{m_1}\|_{\beta,p} \\ &\leq \left[\sum_{j=1}^{m_1+m_2+1} \beta_j^p \right]^{\frac{1}{p}} \leq \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}. \end{aligned}$$

This means that

$$\text{diam}_{\|\cdot\|_{L,\alpha,\beta,p,\bar{p}}} \{e_i\}_i = \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.$$

Now we compute $\lim_m \text{dist}_{\|\cdot\|_{L,\alpha,\beta,p,\bar{p}}}(e_{m+1}, \text{conv}\{e_1, \dots, e_m\})$. Let us take $a^1 + \dots + a^m = 1$, where $0 \leq a^k \leq 1$ for $1 \leq k \leq m$. Then we have

$$e_{m+1} - \sum_{k=1}^m a^k e_k = \left\{ e_j^* \left(e_{m+1} - \sum_{k=1}^m a^k e_k \right) \right\}_j = \{-a^1, \dots, -a^m, 1, 0, \dots\}$$

for all $m \in \mathbb{N}$. But we also have

$$\alpha \left\| e_{m+1} - \sum_{k=1}^m a^k e_k \right\|_{\bar{p}} \leq 1.$$

Hence we get

$$\begin{aligned} \left[\sum_{j=1}^{m+1} \beta_j^p \right]^{\frac{1}{p}} &\leq \left\| e_{m+1} - \sum_{k=1}^m a^k e_k \right\|_{L,\alpha,\beta,p,\bar{p}} \\ &\leq \text{diam}_{\|\cdot\|_{L,\alpha,\beta,p,\bar{p}}} \{e_i\}_i = \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}. \end{aligned}$$

This means that

$$\lim_m \operatorname{dist}_{\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}}(e_{m+1}, \operatorname{conv}\{e_1, \dots, e_m\}) = \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.$$

□

6. The norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ and asymptotic normal structure. We have the following result.

Theorem 6.1. *The Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ fails asymptotic normal structure for $\alpha \leq 4^{-\frac{1}{\tilde{p}}}$.*

Proof. Once again $\{e_i\}_i$ is the standard basis in $l^{\tilde{p}}$. Analogously as in [13] (see also [2]) we put

$$(**) \quad x_n = \begin{cases} (1 - \frac{j}{2^{2i}})e_i + e_{i+1}, & \text{if } n = 2^{2i} + j, \quad j = 1, 2, \dots, 2^{2i} \\ e_{i+1} + \frac{j}{2^{2i+1}}e_{i+2}, & \text{if } n = 2^{2i+1} + j, \quad j = 1, 2, \dots, 2^{2i+1}. \end{cases}$$

and

$$C = \overline{\operatorname{conv}}\{x_n : n = 5, 6, \dots\}.$$

Directly from (**) we get

$$0 = \lim_n \|x_n - x_{n+1}\|_{\tilde{p}} = \lim_n \|x_n - x_{n+1}\|_{L,\alpha,\beta,p,\tilde{p}}$$

and

$$\alpha \operatorname{diam}_{\|\cdot\|_{\tilde{p}}}(C) \leq \alpha 4^{\frac{1}{\tilde{p}}} \leq 1.$$

Next it is easy to see that $|x_n^k - x_m^k| \leq 1$ for every $k, m, n \in \mathbb{N}$ and therefore

$$\|x_n - x_m\|_{L,\alpha,\beta,p,\tilde{p}} \leq \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.$$

On the other hand, for example, if $i_1 + 2 < i_2$, $n_1 = 2^{2i_1} + 1$, $n_2 = 2^{2i_2} + 1 > 2^{2i_1+4} + 1$, then we have

$$\begin{aligned} x_{n_2} - x_{n_1} &= \{x_{n_2}^k - x_{n_1}^k\}_k \\ &= \{0, \dots, 0, x_{n_2}^{i_1} - x_{n_1}^{i_1}, x_{n_2}^{i_1+1} - x_{n_1}^{i_1+1}, 0, \dots, 0, x_{n_2}^{i_2} - x_{n_1}^{i_2}, x_{n_2}^{i_2+1} - x_{n_1}^{i_2+1}, 0, \dots\} \\ &= \left\{0, \dots, 0, \frac{1}{2^{2i_1}} - 1, -1, 0, \dots, 0, 1 - \frac{1}{2^{2i_2}}, 1, 0, \dots\right\}. \end{aligned}$$

and

$$\alpha \|x_{n_2} - x_{n_1}\|_{\tilde{p}} \leq \alpha \operatorname{diam}_{\|\cdot\|_{\tilde{p}}}(C) \leq 1.$$

Thus directly from the definition of the norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ we get

$$\left[\sum_{j=1}^{i_1+i_2+2} \beta_j^p \right]^{\frac{1}{p}} \leq \|x_{n_2} - x_{n_1}\|_{L,\alpha,\beta,p,\tilde{p}} \leq \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.$$

This means that

$$\text{diam}_{L,\alpha,\beta,p,\tilde{p}}(C) = \text{diam}_{\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}} \{x_n\} = \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}}.$$

Similarly, we obtain

$$\left\| \sum_{l=1}^m a_l x_l - x_n \right\|_{L,\alpha,\beta,p,\tilde{p}} \geq \left[\sum_{j=1}^{i+1} \beta_j^p \right]^{\frac{1}{p}}$$

for each convex combination $\sum_{l=1}^m a_l x_l$ and for every sufficiently large n , where i is connected with n by (**). This implies

$$\lim_n \left\| \sum_{l=1}^m a_l x_l - x_n \right\|_{L,\alpha,\beta,p,\tilde{p}} = \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}} = \text{diam}_{\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}}(C).$$

Hence we have

$$\lim_n \|x - x_n\|_{L,\alpha,\beta,p,\tilde{p}} = \text{diam}_{\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}}(C)$$

for each $x \in C$ and therefore the set C lacks asymptotic normal structure. \square

7. The norm $\|\cdot\|_{L,\alpha,\beta,p,\tilde{p}}$ and LUR. Here applying the Smith method (Example 1 in [16]), we arrive at the following theorem.

Theorem 7.1. *The Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ is LUR.*

Proof. Let $x \in l^{\tilde{p}}$ and $\{x_n\}_n \subset l^{\tilde{p}}$ be such that $\|x\|_{L,\alpha,\beta,p,\tilde{p}} = 1$, $\lim_n \|x_n\|_{L,\alpha,\beta,p,\tilde{p}} = 1$ and $\lim_n \|x + x_n\|_{L,\alpha,\beta,p,\tilde{p}} = 2$. Then we also have $\|u(x)\|_{\beta,p} = 1$, $\lim_n \|u(x_n)\|_{\beta,p} = 1$ and $\lim_n \|u(x) + u(x_n)\|_{\beta,p} = 2$. Applying local uniform convexity (LUR) of $(c_0, \|\cdot\|_{\beta,p})$, we immediately obtain the strong convergence of the sequence $\{u(x_n)\}$ to $u(x)$ in the norm $\|\cdot\|_{\beta,p}$. But we have

$$\begin{aligned} \beta_1 \|u(x) - u(x_n)\|_{c_0} &\leq \|u(x) - u(x_n)\|_{\beta,p} \\ &\leq \left[\sum_{j=1}^{\infty} \beta_j^p \right]^{\frac{1}{p}} \|u(x) - u(x_n)\|_{c_0} \end{aligned}$$

and

$$\|u(x) - u(x_n)\|_{c_0} = \max\{\alpha \| \|x\|_{\tilde{p}} - \|x_n\|_{\tilde{p}} \}, |x^1 - x_n^1|, |x^2 - x_n^2|, \dots \}.$$

This implies that $\lim_n \|x_n\|_{\tilde{p}} = \|x\|_{\tilde{p}}$ and $\lim_n x_n^i = x^i$ for $i = 1, 2, \dots$. Hence the sequence $\{x_n\}$ tends weakly to x . Finally, by the Kadec–Klee property of $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ we have $\lim_n x_n = x$ in $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ and therefore in $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$, too. \square

8. Diametrically complete sets with empty interior in the space $(l^{\bar{p}}, \|\cdot\|_{L,\alpha,\beta,p,\bar{p}})$. Using the properties of the norm $\|\cdot\|_{L,\alpha,\beta,p,\bar{p}}$ and the Maluta–Papini Theorem, we obtain

Theorem 8.1. *For $\alpha \leq 2^{-\frac{1}{\bar{p}}}$ the Banach space $(l^{\bar{p}}, \|\cdot\|_{L,\alpha,\beta,p,\bar{p}})$ is LUR and contains a diametrically complete set whose interior is empty.*

Proof. It is a simple consequence of Theorems 2.4, 4.1, 5.1 and 7.1. \square

9. The weak fixed point property. We begin with the following definition.

Definition 9.1. Let $(X, \|\cdot\|)$ be a Banach space with a Schauder basis $\{e_i\}_i$. We say that a Schauder basis $\{e_i\}_i$ is unconditional if whenever the series $\sum_{i=1}^{\infty} x^i e_i$ converges, it converges unconditionally, i.e., $\sum_{i=1}^{\infty} x^{\tilde{\sigma}(i)} e_{\tilde{\sigma}(i)}$ converges for every permutation $\tilde{\sigma}$ of \mathbb{N} .

The following theorem shows a few equivalent definitions of unconditionality of a Schauder basis (see for example [10] and [15]).

Theorem 9.2. *Let $(X, \|\cdot\|)$ be a Banach space with a Schauder basis $\{e_i\}_i$. The following statements are equivalent:*

- (1) *the basis $\{e_i\}_i$ is an unconditional basis;*
- (2) *for every choice of signs $\{\varepsilon_i\}_i$ (i.e. $\varepsilon_i = \pm 1$) $\sum_{i=1}^{\infty} \varepsilon_i x^i e_i$ converges whenever $\sum_{i=1}^{\infty} x^i e_i$ converges;*
- (3) *for every convergent series $\sum_{i=1}^{\infty} x^i e_i$ and for every sequence of scalars $\{b^i\}_i$ such that $|b^i| \leq |x^i|$ for all i the series $\sum_{i=1}^{\infty} b^i e_i$ converges.*

It is known that if $\{e_i\}_i$ is an unconditional Schauder basis in a Banach space $(X, \|\cdot\|)$, then

$$\sup \left\{ \left\| \sum_{i=1}^{\infty} \varepsilon_i x^i e_i \right\| : \left\| \sum_{i=1}^{\infty} x^i e_i \right\| = 1 \text{ and } \varepsilon_i = \pm 1 \right\}$$

is finite (see for example [10] and [15]). Hence we state

Definition 9.3. If a Schauder basis $\{e_i\}_i$ is an unconditional basis in a Banach space $(X, \|\cdot\|)$, then the number

$$K := \sup \left\{ \left\| \sum_{i=1}^{\infty} \varepsilon_i x^i e_i \right\| : \left\| \sum_{i=1}^{\infty} x^i e_i \right\| = 1 \text{ and } \varepsilon_i = \pm 1 \right\}$$

is called the unconditional constant of $\{e_i\}_i$. If this constant K is equal to 1, then we say that $\{e_i\}_i$ is a 1-unconditional basis.

Next we recall the definition of the weak fixed point property.

Definition 9.4. Let $(X, \|\cdot\|)$ be a Banach space and let C be a weakly compact and convex subset of X . We say that C has the fixed point property if each nonexpansive mappings $T : C \rightarrow C$ (i.e. $\|Tx_1 - Tx_2\| \leq \|x_1 - x_2\|$ for every $x_1, x_2 \in C$) has a fixed point.

If each weakly compact and convex subset C of X has the fixed point property, then we say that a Banach space $(X, \|\cdot\|)$ has the weak fixed point property.

The following theorem is generally known.

Theorem 9.5 ([4]). *Let $(X, \|\cdot\|)$ be a Banach space, $C \subset X$ be weakly compact and convex and let $T : C \rightarrow C$ be a nonexpansive mapping. Then there exists a separable subspace $X_1 \subset X$ such that the set $C_1 = C \cap X_1$ is a weakly compact, convex and T -invariant, i.e. $T(C_1) \subset C_1$.*

In 1965, W. A. Kirk [7] published his famous fixed point theorem.

Theorem 9.6. *Let C be a nonempty, weakly compact, convex subset of a Banach space $(X, \|\cdot\|)$, and suppose C has normal structure. Then each nonexpansive mapping $T : C \rightarrow C$ has a fixed point.*

Next in 1981, J. B. Baillon and R. Schöneberg [2] extended Kirk's Theorem, using asymptotic normal structure.

Theorem 9.7. *Every reflexive Banach space with asymptotic normal structure has the weak fixed point property for nonexpansive mappings.*

Observe that by Theorems 7.6 and 8.2 from [3] we are not able to apply the above fixed point theorems to the Banach space $(c_0, \|\cdot\|_{\beta,p})$.

However, the following theorem, which is due to P.-K. Lin ([9]), is valid.

Theorem 9.8. *Each Banach space $(X, \|\cdot\|)$ with a 1-unconditional Schauder basis $\{e_i\}_i$ has the weak fixed point property.*

Directly from the definitions of a 1-unconditional Schauder basis and the norm $\|\cdot\|_{\beta,p}$ we get

Theorem 9.9. *Let $\{e_i\}_i$ be a standard basis in $c_0 = c_0(\mathbb{N})$. Then $\{e_i\}_i$ is a 1-unconditional Schauder basis in $(c_0, \|\cdot\|_{\beta,p})$.*

Thus we are ready to prove

Theorem 9.10. *The Banach space $(c_0(\Gamma), \|\cdot\|_{\beta,p})$ has the weak fixed point property.*

Proof. By Theorem 9.5 we can assume that $\Gamma = \mathbb{N}$ and $c_0(\Gamma) = c_0$. Since by Theorem 9.9 the standard basis $\{e_i\}$ in c_0 is a 1-unconditional Schauder basis in $(c_0, \|\cdot\|_{\beta,p})$, we can apply Theorem 9.8 to get our result. \square

Next, if $\alpha \leq 4^{-\frac{1}{\tilde{p}}}$, then by Theorem 5.1 and 6.1 we are not able to apply Theorems 9.6 and 9.7 to the Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$. But directly from the definitions of a 1-unconditional Schauder basis and the norm $\|\cdot\|_{\beta,p}$, we obtain

Theorem 9.11. *Let $\{e_i\}_i$ be a standard basis in $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$. Then $\{e_i\}_i$ is a 1-unconditional Schauder basis in $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$.*

So we are ready to prove

Theorem 9.12. *The Banach space $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$ has the weak fixed point property.*

Proof. Since by Theorem 9.11 the standard basis $\{e_i\}$ in $(l^{\tilde{p}}, \|\cdot\|_{\tilde{p}})$ is a 1-unconditional Schauder basis in $(l^{\tilde{p}}, \|\cdot\|_{L,\alpha,\beta,p,\tilde{p}})$, the Lin Theorem implies our result. \square

Acknowledgments. These results have been partially achieved within the framework of the STREVCOMS Project No. 612669 with funding from the IRSES Scheme of the FP7 Programme of the European Union (the first author).

REFERENCES

- [1] Ayerbe Toledano, J. M., Domínguez Benavides, T., López Acedo, G., *Measures of Noncompactness in Metric Fixed Point Theory*, Birkhäuser, 1997.
- [2] Baillon, J.-B., Schöneberg, R., *Asymptotic normal structure and fixed points of non-expansive mappings*, Proc. Amer. Math. Soc. **81** (1981), 257–264.
- [3] Budzyńska, M., Grzesik, A., Kot, M., *The generalized Day norm. Part I. Properties*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **71** (2) (2017), 33–49.
- [4] Goebel, K., Kirk, W. A., *Topics in Metric Fixed Point Theory*, Cambridge University Press, 1990.
- [5] Holmes, R. B., *Geometric Functional Analysis and Its Applications*, Springer, 1975.
- [6] Kadec, M. I., *On the connection between weak and strong convergence*, Dopovidi Akad. Nauk Ukrain. RSR **9** (1959), 949–952.
- [7] Kirk, W. A., *A fixed point theorem for mappings which do not increase distances*, Amer. Math. Monthly **72** (1965), 1004–1006.
- [8] Klee, V., *Mappings into normed linear spaces*, Fund. Math. **49** (1960/1961), 25–34.
- [9] Lin, P.-K., *Unconditional bases and fixed points of nonexpansive mappings*, Pacific J. Math. **116** (1985), 69–76.
- [10] Lindenstrauss, J., Tzafriri, L., *Classical Banach Spaces I and II*, Springer, 1977.
- [11] Maluta, E., *A diametrically complete set with empty interior in a reflexive LUR space*, J. Nonlinear Conv. Anal. **18** (2017), 105–111.
- [12] Maluta, E., Papini, P. L., *Diametrically complete sets and normal structure*, J. Math. Anal. Appl. **424** (2015), 1335–1347.
- [13] Mariadoss, S. A., Soardi, P. M., *A remark on asymptotic normal structure in Banach spaces*, Rend. Sem. Mat. Univ. Politec. Torino **44** (1986), 393–395.
- [14] Moreno, J. P., Papini, P. L., Phelps, R. R., *Diametrically maximal and constant width sets in Banach spaces*, Canad. J. Math. **58** (2006), 820–842.
- [15] Singer, I., *Bases in Banach Spaces I*, Springer, 1970.

- [16] Smith, M. A., *Some examples concerning rotundity in Banach spaces*, Math. Ann. **233** (1978), 155–161.
- [17] Smith, M. A., Turett, B., *A reflexive LUR Banach space that lacks normal structure*, Canad. Math. Bull. **28** (1985), 492–494.

Monika Budzyńska
Institute of Mathematics
Maria Curie-Skłodowska University
20-031 Lublin
Poland
e-mail: monikab1@hektor.umcs.lublin.pl

Aleksandra Grzesik
Faculty of Mathematics
and Applied Physics
Rzeszow University of Technology
35-959 Rzeszów, Poland
e-mail: a.grzesik22@gmail.com

Mariola Kot
Faculty of Mathematics and Applied Physics
Rzeszow University of Technology
35-959 Rzeszów, Poland
e-mail: m.kot@prz.edu.pl

Received October 11, 2017