

JACEK KOBA

Forest Management and Geodesy Bureau, Branch in Lublin
PO-20-352 Lublin, ul. Startowa 11, jacek.koba@lublin.buligl.pl
(PhD student in the Department of Botany, Jan Kochanowski University in Kielce)

Alien and invasive plant species in plant communities of floodplain forests of the Małopolska Upland

Obce oraz inwazyjne gatunki roślin w zbiorowiskach lasów łągowych Wyżyny Małopolskiej

SUMMARY

During the field studies on the floodplain forests of the Małopolska Upland carried out in 2009–2013, the author tried to determine which species of alien and invasive plants could be a threat to the stability of these vegetation communities. Efforts were also made to answer the question, which floodplain forests are particularly vulnerable to the penetration of alien species and in which layers of the forest plant community are the best conditions for the development of the individual alien plant species. During the research, a total number of 344 relèves were made, of which 90 recorded the presence of alien species. The most common were: *Impatiens parviflora*, *Acer negundo* and *Padus serotina*. The highest percentage of relèves involving alien species was found in the riverside floodplain forests of *Salicetum albo-fragilis* and *Populetum albae*.

Keywords: invasive alien plant species, floodplain forests, Małopolska Upland

STRESZCZENIE

Podczas prac terenowych poświęconych szacie roślinnej lasów łągowych Wyżyny Małopolskiej przeprowadzonych w latach 2009–2013, autor podjął próbę określenia, jakie gatunki roślin obcych oraz inwazyjnych mogą stanowić zagrożenie dla tych zbiorowisk roślinnych. Starano się również dać odpowiedź na pytanie, które zespoły lasów łągowych są szczególnie podatne na wnikanie gatunków obcych oraz w jakich warstwach leśnego zbiorowiska roślinnego znajdują warunki do rozwoju poszczególne obce gatunki roślin. W ramach prac badawczych wykonano 344 zdjęcia fitosocjologiczne, spośród których w obrębie 90 powierzchni zanotowano występowanie gatunków obcych. Do najczęściej spotykanych należały: *Impatiens parviflora*, *Acer negundo*

oraz *Padus serotina*. Największy odsetek zdjęć z udziałem gatunków obcych stwierdzono w fitocenozach nadrzecznych łąg wierzbowego *Salicetum albo-fragilis* oraz nadrzecznych łąg topolowego *Populetum albae*.

Słowa kluczowe: obce inwazyjne gatunki roślin, lasy łąkowe, Wyżyna Małopolska

INTRODUCTION

Invasions of alien species is now considered one of the most important problems of nature conservation on a global scale (11), which also has an increasing impact on the economic function of forests (17). Based on field studies on the floodplain forests of the Małopolska Upland, the presented work is an attempt to determine which species of alien and invasive plants could be a threat to the stability of these vegetation communities. Efforts were also made to answer the question, which communities of the floodplain forests are particularly vulnerable to the penetration of alien species and in which layers of the forest plant community create the conditions for the development of individual alien plant species.

Floodplain forests are a specific type of vegetation associated primarily with the valleys of watercourses. These communities were the subject of several monographs in Poland and among them are mentioned works by: Matuszkiewicz et al. (24), Faliński (9), Kasproicz and Wojterska (16), Matuszkiewicz (22, 23). The floodplain forests of the Małopolska Upland areas currently do not have a collective monography, and information about them can be found in among others works by: Barański et al. (1), Fabijanowski and Zarzycki (7, 8), Głazek (12, 13, 14), Bróż, (3), Różański (33), Hereźniak (15) and Przemyski (29, 30, 31). There is also no separate research dedicated to participation and role of invasive alien species as a component of floodplain forests.

STUDY AREA, MATERIALS AND METHODS

The field studies were carried out between 2009 and 2013. The works included the entire area of the Małopolska Upland within the borders set by Kondracki (20). The research included, among other things, the valleys of the main rivers flowing through the area such as: the Pilica, Nida, Kamienna, Warta and Vistula.

In total, 344 relèves were made. The relèves were made for the characterisation of plant communities in floodplain forests. Next, an analysis on the prevalence of alien and invasive species was carried out on the entire phytosociological database.

A description of vegetation has been prepared in accordance with the methodology presented by Braun-Blanquet (2). The selection of research areas was made after earlier analysis of forest site types, soil and water conditions in the Małopolska Upland forests, using documents such as forest management plans, and available layers of digital maps [GIS] containing information about the diversity of vegetation. The nomenclature of species was accepted following Mirek et al. (25) and forest communities – Matuszkiewicz (23). The invasive alien species were considered according to the proposal of Tokarska-Guzik et al. (37).

RESULTS

As a result of phytosociological research, five floodplain forest associations were determined: *Ficario-Ulmetum minoris* Knapp 1942 em. J. Mat. 1976, *Fra-*

xino-Alnetum W. Mat. 1952, *Carici remotae-Fraxinetum* Koch 1926, *Salicetum albo-fragilis* Tx. 1955 and *Populetum albae* Br.-Bl. 1931. In addition, the secondary forest communities (*Alnus glutinosa-Rubus idaeus*, *Alnus glutinosa-Urtica dioica*) related to floodplain forests (26, 27), as were indicated.

In all the floodplain forest communities, the presence of alien and invasive species has been identified. Among the 344 reléves, 90 recorded the presence of alien species (Fig. 1), which is 26% of all localities.




Fig. 1. Location of the study sites and sites of alien and invasive species in floodplain forests comprising in the boundary of the Malopolska Upland within the ATPOL grid [10 km x 10 km].

In total, 15 alien and invasive species of vascular plants were found (Tab. 1). Among the described plants are the following forms: nine species of trees, one shrub species, one species of climbers, two species of perennials and two species of annual plants. The majority of alien species habitats were near the banks of rivers and watercourses. The occurrence of alien species was quite often reported also within small forest patches separated from the major forest areas.

Tab. 1. Alien and invasive species in communities of floodplain forests of the Malopolska Upland.

No.	Latin name of species	Family	Geographical -historical group	Current status	Number of sites [in relèves]
1.	<i>Acer negundo</i> L.	<i>Aceraceae</i>	Kn	invasive	16
2.	<i>Aesculus hippocastanum</i> L.	<i>Hippocastanaceae</i>	Kn	-----	3
3.	<i>Fraxinus pennsylvanica</i> Marshall	<i>Oleaceae</i>	Kn	invasive	1
4.	<i>Impatiens glandulifera</i> Royle	<i>Balsaminaceae</i>	Kn	invasive	2
5.	<i>Impatiens parviflora</i> DC.	<i>Balsaminaceae</i>	Kn	invasive	53
6.	<i>Juglans regia</i> L.	<i>Juglandaceae</i>	Kn	-----	2
7.	<i>Malus domestica</i> Borkh.	<i>Rosaceae</i>	Kn	-----	2
8.	<i>Padus serotina</i> [Ehrh]. Borkh.	<i>Rosaceae</i>	Kn	invasive	10
9.	<i>Parthenocissus inserta</i> [A. Kern] Fritsch	<i>Vitaceae</i>	Kn	invasive	2
10.	<i>Pyrus communis</i> L.	<i>Rosaceae</i>	Kn	-----	4
11.	<i>Quercus rubra</i> L.	<i>Fagaceae</i>	Kn	invasive	2
12.	<i>Robinia pseudacacia</i> L.	<i>Fabaceae</i>	Kn	invasive	2
13.	<i>Solidago canadensis</i> L.	<i>Asteraceae</i>	Kn	invasive	5
14.	<i>Solidago gigantea</i> Aiton	<i>Asteraceae</i>	Kn	invasive	8
15.	<i>Symphoricarpos albus</i> [L.] S.F. Blake	<i>Caprifoliaceae</i>	Kn	-----	1

Kn – kenophyte

The most common invasive species was *Impatiens parviflora*, which appeared in 53 relèves. The species also relatively frequent were: *Acer negundo*, *Padus serotina*, *Solidago gigantea* and *Solidago canadensis*.

Impatiens parviflora was most often found species in a community of *Fra-xino-Alnetum* [28 localities] and *Ficario-Ulmetum minoris* [18 localities]. The second, in terms of the number of localities is *Acer negundo*, occurring mainly in *Salicetum albo-fragilis* and *Populetum albae*. In addition, *Acer negundo* was found quite frequently within a community of *Ficario-Ulmetum minoris*.

Taking into account the percentage of the area affected by alien species in different communities (Tab. 2), it can be observed that the most common finding alien and invasive species [92% – 100% of the relèves in phytocoenosis] relate to

Tab. 2. Relation of occurrence of alien and invasive species and the communities of floodplain forests [number of sites in the associations].

No.	Invasive alien species	<i>Fraxino-Alnetum</i>	<i>Carici remotae-Fraxinetum</i>	<i>Ficario-Ulmetum minoris</i>	<i>Salicetum albo-fragilis</i>	<i>Populetum albae</i>	The secondary communities
1.	<i>Acer negundo</i> L.	1	-	7	2	6	-
2.	<i>Aesculus hippocastanum</i> L.	1	-	2	-	-	-
3.	<i>Fraxinus pennsylvanica</i> Marshall	1	-	-	-	-	-
4.	<i>Impatiens glandulifera</i> Royle	-	-	-	-	-	2
5.	<i>Impatiens parviflora</i> DC.	28	3	18	-	4	-
6.	<i>Juglans regia</i> L.	-	-	-	-	2	-
7.	<i>Malus domestica</i> Borkh.	2	-	-	-	-	-
8.	<i>Padus serotina</i> [Ehrh.], Borkh.	7	-	3	-	-	-
9.	<i>Parthenocissus inserta</i> [A. Kern] Fritsch	-	-	2	-	-	-
10.	<i>Pyrus communis</i> L.	3	-	1	-	-	-
11.	<i>Quercus rubra</i> L.	1	1	-	-	-	-
12.	<i>Robinia pseudacacia</i> L.	-	-	1	-	1	-
13.	<i>Solidago canadensis</i> L.	1	-	3	-	-	1
14.	<i>Solidago gigantea</i> Aiton	2	1	-	2	3	-
15.	<i>Symphoricarpos albus</i> [L.] S.F. Blake	1	-	-	-	-	-
Total number of relèves in association:		246	12	62	3	12	9
Number of relèves with invasive alien species in association:		41	5	27	3	11	3
Percentage (%) of relèves with invasive alien species in association:		17	42	44	100	92	33

communities *Salicetum albo-fragilis* and *Populetum albae*. The average percentage [33% – 44% of all relèves] falls into the communities of *Ficario-Ulmetum minoris*, *Carici remotae-Fraxinetum* and secondary forest communities. The most frequent community in this study of *Fraxino-Alnetum*, is the subject of invasions, relatively to say the least, because the percentage of relèves with the participation of alien species is only about 17% in the total number of the studied locations.

Taking into account the presence and coverage of various alien species of trees and shrubs in forest layers – it should be noted that the highest degree of expansiveness had *Acer negundo* (Tab. 3). This species is encountered in all layers, from the herb one, to the canopy and reaching a maximum coverage of 20%. Another tree species with a relatively high degree of expansiveness is *Padus serotina*, and is present in herb and shrubs layers with a maximum coverage up to 30%. It should be noted, however, that this species did not occur in the tree layer. In addition, low levels of coverage listed in the herb layer may indicate a difficult sowing of *Padus serotina* in floodplain forests, which probably is connected with inadequate habitat conditions for this species.

Among herbaceous species, the greatest cover is up to 60% and is achieved locally by *Impatiens glandulifera*. This plant, however, was recorded in only two relèves. Definitely more frequent in the herb layer are alien invasive species, such as *Impatiens parviflora* and *Solidago canadensis*, which reach up to 40% coverage. These species are extremely affected by the physiognomy of the herb layer for floodplain forest communities. However, there is no evidence that these plants form within the surveyed area of clear single-species aggregations.

DISCUSSION

Research carried out within the flood forest communities of the Małopolska Upland, confirms the thesis of other authors (10, 21, 39), which indicate riverine habitats as particularly vulnerable to invasions of alien species. The spread of these species in the riverine alluvial forests is often associated with the thinning of plant communities, as well as with the presence of many routes, which was observed by the author in the valleys of the Warta, Pilica, Nida and Opatówka.

Studies conducted in the Małopolska Upland indicate, like the other works (18), that the *Fraxino-Alnetum* as a community is relatively mostly resistant to the penetration of alien and invasive species. The factors limiting processes of invasion are undoubtedly strong shading, variable moisture and less human impact caused by the unavailability of these forests. This association has the relatively smallest degree of degradation in comparison with other communities of floodplain forests. It can therefore be assumed, that a stable community of *Fraxino-Alnetum*, has a high degree of resistance to the expansion of alien species.

Tab. 3. Alien and invasive species in layers of the floodplain forest communities.

No.	Invasive alien species in layers	Layers	Cover min. %	Cover max. %	Cover aver. %
Trees and shrubs					
1	<i>Acer negundo</i> L.	A1	1	20	9
		A2	1	20	11
		B	1	20	7
		C	1	5	2
2	<i>Aesculus hippocastanum</i> L.	A2	-	1*	-
		B	-	1*	-
		C	-	1*	-
3	<i>Fraxinus pennsylvanica</i> Marshall	A2	-	20*	-
4	<i>Juglans regia</i> L.	A2	-	1*	-
		B	-	1*	-
5	<i>Malus domestica</i> Borkh.	A2	-	1**	-
6	<i>Padus serotina</i> [Ehrh.] Borkh.	B	1	30	5
		C	1	1	1
7	<i>Pyrus communis</i> L.	B	1	1	1
		C	-	1*	-
8	<i>Quercus rubra</i> L.	B	-	1**	-
		C	-	1*	-
9	<i>Robinia pseudacacia</i> L.	B	-	1*	-
10	<i>Symphoricarpos albus</i> [L.] S.F. Blake	B	-	15*	-
Herbaceous plants, climbers					
11	<i>Impatiens glandulifera</i> Royle	C	15*	60*	-
12	<i>Impatiens parviflora</i> DC.	C	1	40	9
13	<i>Parthenocissus inserta</i> [A. Kern] Fritsch	C	1*	15*	-
14	<i>Solidago canadensis</i> L.	C	1	40	13
15	<i>Solidago gigantea</i> Aiton	C	1	15	7

* 1 stand; ** 2 stands

A species, which is a significant threat to the natural structure of floodplain forest communities, is *Acer negundo*, and can be found in the Małopolska Upland habitat conditions similar to those that occur in the natural range in North America (4). Its presence in all layers of the forest community is noted especially in the riverine alluvial forests. This species, in extreme cases, changes the physiognomy of the landscape of river valleys, as observed by the author in the Opatówka, Szreniawa and Nida valleys. Another invasive species that can significantly modify the structure of floodplain forests is *Impatiens parviflora*. Together with *Acer negundo*, this species is potentially the greatest threat to the floodplain forests phytocoenoses of the Małopolska Upland.

All alien and invasive species found in floodplain forests of the Małopolska Upland were earlier reported by other authors from similar habitats (5, 6, 18, 19, 21, 28, 32, 34–36, 38). During the research conducted in the Małopolska Upland, no such invasive species as *Echinocystis lobata* and *Reynoutria spp* were found. This is in contradiction to facts being quite often reported from floodplain phytocoenoses by other authors (5, 18, 34). Limited possibilities for the spread of these taxons in the floodplain forests presented in this paper confirm the thesis put forward by the above authors, that these species spread intensively only in open areas or within heavily overexposed stands of forests. Their penetration into relatively dense, semi-natural phytocoenoses examined by the author is probably difficult.

CONCLUSIONS

1. The presence of alien and invasive species of vascular plants was confirmed in all types of floodplain forests in the Małopolska Upland.
2. Most susceptible to the processes of penetration of alien species are the following communities of riverside forests: *Salicetum albo-fragilis*, *Populetum albae*, and *Ficario-Ulmetum minoris*.
3. Alien species encountered most often in floodplain forests are: *Impatiens parviflora*, *Acer negundo*, and *Padus serotina*.
4. The greatest threat to the natural structure of floodplain forests are *Impatiens parviflora* and *Acer negundo* which can significantly disrupt the physiognomy of plant communities.
5. Phytocenosis most resistant to the penetration of alien species is *Fraxino-Alnetum*.

REFERENCES

1. Barański S., Bober L., Adamczyk B., Fabijanowski J., Zarzycki K., Kuc M., Nowak J. 1960. The forest reserve Świnia Góra. State Council for Conservation of Nature. Poland. Warsaw–Kraków. 16: 3–23.

2. Braun-Blanquet J. 1928, 1964. Pflanzensociologie. Springer, Wien–New York.
3. Bróz E. 1985. Szata roślinna rezerwatu Czarny Las w Świętokrzyskim Parku Narodowym. Rocz. Świętokrz. 12: 99–123.
4. Carter Johnson W., Burghess L.R., Keammerer R. 1976. Forest overstory vegetation and environment on the Missouri river floodplain in North Dakota.
5. Dajdok Z., Kącki Z. 2003. Kenophytes of the Odra riversides, in: A. Zając, M. Zając, B. Zemanek [eds.], Phytogeographical Problems of Synanthropic Plants, Institute of Botany Jagiellonian University, Cracow, pp. 131–136.
6. Danielewicz W. 1993. Obce gatunki drzew i krzewów w dolinie Warty. 1. Klon jesionolistny (*Acer negundo* L.). Prace Kom. Nauk Roln. i Kom. Nauk Leśn. PTPN 76, 31–37.
7. Fabijanowski J., Zarzycki K. 1965. Roślinność rezerwatu leśnego Świnia Góra w Górach Świętokrzyskich. Acta Agr. et Silv., ser. leśna. Vol. 5: 63–103.
8. Fabijanowski J., Zarzycki K. 1967. Wody gruntowe w zbiorowiskach leśnych Nadleśnictwa Bliżyn [Góry Świętokrzyskie]. Acta Agr. Silv., ser. Silv. 7: 3–39.
9. Faliński J.B. 1961. Zbiorowiska łąkowe Krainy Wielkich Jezior Mazurskich – mscr. Praca doktorska w Zakł. Fitosocjol. Uniw. Warsz.
10. Faliński J.B. 2000. Rzeczne wędrówki roślin. Rzeki: Kultura – Cywilizacja – Historia. 9, pp. 143–186.
11. Genovesi P., Shine C. 2004. European strategy on invasive alien species. Nature and Environment 137. Council of Europe Publishing, Strasbourg, France, pp. 1–67. http://www.coe.int/t/dg4/cultureheritage/conventions/Bern/T-PVS/sc24_inf01_en.pdf.
12. Głazek T. 1973. Zespoły leśne północno-wschodniego i wschodniego przedpola Gór Świętokrzyskich. Monogr. Bot. 38: 1–158.
13. Głazek T. 1985. Szata roślinna wybranych powierzchni obszaru Gór Świętokrzyskich i terenów przyległych na tle warunków siedliskowych. Fragm. Faun. 29 [11]: 153–234.
14. Głazek T., Wolak J. 1991. Zbiorowiska roślinne Świętokrzyskiego Parku Narodowego i jego strefy ochronnej. Monographiae Botanicae 72: 3–108.
15. Hereźniak J. 1993. Stosunki geobotaniczno-leśne północnej części Wyżyny Śląsko-Krakowskiej na tle zróżnicowania i przemian środowiska. Monogr. Bot. 75: 5–364.
16. Kasprówicz M., Wojterska M. 1988. Olesy, łągi olszowe i wiązowe oraz ich formy degeneracyjne w okolicach Konina. PTPN, Prace Kom. Biol. 70: 170–163.
17. Kettunen M., Genovesi P., Gollasch S., Pagad S., Starfinger U., ten Brink P., Shine C. 2009. Technical support to EU strategy on invasive species [IAS] – Assessment of the impacts of IAS in Europe and the EU [Final draft report for the European Commission]. Institute for European Environmental Policy [IEEP], Brussels, Belgium, pp. 1–131.
18. Kołaczowska E. 2010. Obce inwazyjne gatunki roślin w krajobrazie dolin Świdra i Rządzy. Prace Komisji Krajobrazu Kulturowego nr 13. Sosnowiec.
19. Kołaczowska E. 2012. Siedliskowo-ekologiczne wzorce występowania obcych inwazyjnych gatunków roślin w dolinach małych rzek wschodniego Mazowsza. Wybrane aspekty. Studia i Materiały CEPL w Rogowie. Vol. 14. Zeszyt 33/4/2012.
20. Kondracki J. 2002. Geografia regionalna Polski. PWN, Warszawa.
21. Kucharczyk M., Krawczyk R. 2004. Kenophytes as River corridor plants in the Vistula River and the San River Valleys. Teka Kom. Ochr. Kształt. Rod. Przyr. 1: 110–115.
22. Matuszkiewicz J.M. 1976. Przegląd fitosocjologiczny zbiorowisk leśnych Polski, cz. 3. Lasy i zarośla łąkowe. Phytocenosis 5 [1]: 3–66.
23. Matuszkiewicz J.M. 2008. Zespoły leśne Polski. Wydawnictwo Naukowe, PWN, Warszawa.
24. Matuszkiewicz W., Traczyk H., Traczyk T. 1958. Materiały do fitosocjologicznej systematyki zespołów olsowych w Polsce. Acta Soc. Bot. Pol., 27.1. Warszawa.

25. Mirek Z., Piękoś-Mirkowa H., Zając A., Zając M. 2002. Krytyczna lista roślin naczyniowych Polski. *Pol. Bot. Studies. Guidebook* 15: 1–442.
26. Olaczek R. 1972. Formy antropogenicznej degeneracji leśnych zbiorowisk roślinnych w krajo-brazie rolniczym Polski niżowej. Łódź, Wyd. UŁ, pp. 1–70.
27. Olaczek R. 1974. Kierunki degeneracji fitocenozy leśnych i metody ich badania. *Phytocoenosis*. 3(3/4): 179–190.
28. Orzechowski M. 2007. Przemiany zbiorowisk roślinnych Puszczy Kozienickiej od czasu badań Ryszarda Zaręby, in: Matuszkiewicz J.M. [ed.] *Geobotaniczne rozpoznanie tendencji rozwojowych zbiorowisk leśnych w wybranych regionach Polski*. Polska Akademia Nauk. Instytut Geografii i Przestrzennego Zagospodarowania im. Stanisława Leszczyckiego. Mon. 8, Warszawa.
29. Przemyski A. 1998. Zaslugujące na ochronę obszary leśne okolic Staszowa w Ziemi Sandomierskiej, in: Puszkarski T. [ed.] *Osobliwości przyrody Ziemi Sandomierskiej. Materiały z sympozjum Sandomierz 16 październik 1998*. Tow. Nauk. Sand, pp. 39–47.
30. Przemyski A., Polinowska K. 2001. Szata roślinna rezerwatu Białe Ługi, in: Żurek S. [ed.] *Rezerwat torfowiskowy „Białe Ługi”*. Wyd. Homini, pp. 133–178.
31. Przemyski A. 2008. Mapa roślinności rzeczynowej i przemiany zbiorowisk w rezerwacie „Białe Ługi”, in: S. Żurek [ed.] *Torfowiska gór, wyżyn i nizin*. Wyd. Uniw. Hum.-Przyr., Kielce, pp. 11–115.
32. Pyšek P., Prach K. 1993. Plant invasions and the role of riparian habitats: a comparison of four species alien to central Europe. *J. Biogeogr.* 20: 413–420.
33. Różański W. 1987. Zróżnicowanie i zachowanie zbiorowisk leśnych Niecki Nidziańskiej. *Studia Ośr. Dok. Fizjogr.* 15: 209–283.
34. Śliwiński M. 2008. Wybrane an tropofity brzegów Bystrzycy na odcinku Krasków–Jarnołtów. *Acta Botanica Silesiaca* 3: 121–136.
35. Śliwiński M., Dajdok Z. 2010. What have we learnt about the invasive plants of the south-western Poland, from the Polish Ecological Club Project? *Acta Botanica Silesiaca* 5: 27–42.
36. Tokarska-Guzik B., 2005. The Establishment and Spread of Alien Plant Species [Kenophytes] in the Flora of Poland, Wydawnictwo Uniwersytetu Śląskiego, Katowice, pp. 1–192.
37. Tokarska-Guzik B., Dajdok Z., Zając M., Zając A., Urbisz A., Danielewicz W., Hołyński Cz. 2012. Rośliny obcego pochodzenia ze szczególnym uwzględnieniem gatunków inwazyjnych. Generalna Dyrekcja Ochrony Środowiska. Warszawa, pp. 1–196.
38. Tylakowski T. 2010. Przekształcenia w składzie dendroflory w dolinie środkowej Warty. *Acta Sci. Pol., Administratio Locorum* 9 (3) 2010, 117–124.
39. Więcław H., Pieńkowski P. 2007. Inwazje gatunków obcych jako element zagrożenia małych dolin rzecznych, in: Myga-Piątek U. [ed.] *Doliny rzeczne: Przyroda Krajobraz – Człowiek*. Prace Komisji Krajobrazu Kulturowego PTG 7, pp. 159–166.